the joints binodose, with whorls of long, white hairs. The abdomen is polished black, ovate, attached to the thorax by a long, punctate petiole.

Described from one & specimen from Tepic.

Perilampus Latreille.

32. Perilampus triangularis Say.

LeConte Ed. Say's Works, i, p. 381.

Six specimens from San Lazaro.

ORASEMA Cameron.

- 33. Orasema viridis sp. n.
- Q.—Length 3 mm. Head, thorax, scape of antennæ, femora and petiole of abdomen, all metallic green; flagellum black; mandibles rufous, the right with two teeth, the left with only one within; tibiæ and tarsi brownish-yellow; abdomen æneous black; wings hyaline, the veins dark brown.

Described from one & specimen from Tepic.

METAPON Walker.

- 34. METAPON MEXICANUM sp. n.
- Q.—Length 2 mm. Head and thorax æneous black, closely punctate; mandibles rufous; scape, pedicel and legs, except the coxæ and anterior and middle femora, ferruginous; anterior and middle femora metallic brown; hind coxæ blue. Abdomen pointed ovate, polished black, as long as the thorax and keeled beneath. Flagellum incrassated toward tip, the joints, after the first, increasing in width and wider than long. Wings hyaline, the veins brown, the marginal vein twice as long as the stigmal. The fourth abdominal segment is not quite half as long as the third, the fifth only a little shorter than the third, while the sixth and seventh are a little longer.

Described from one 9 specimen from Tepic.

2D SER., Vol. V.

(36)

September 7, 1895.