

33. *Subfam. Eucharissinae m.*

SACCHARISSA

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 37.

Etymol. obscura.

synon.: Eucharis Walker.*contingens* (Walk.) Kby. — ♂ — As.: Borneo, Sarawak.

Eucharis contingens Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 378, ♂.

Saccharissa contingens W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 37;

[T. 1 F. 6.]

EUCCHARISSA

Westwood, Trans. Entom. Soc. London 1868 Proc. p. XXXVI.

*εὐχαρῖς, gratus.**cuprifrons* Westw. — ♀ — Afr.: Caffraria.

Eucharissa cuprifrons Westwood, Thesaur. entom. Oxon. 1874 p. 144 n. 4, ♀.

natalica Westw. — Afr.: Natal.

Eucharissa Natalica Westwood, Trans. Entom. Soc. London 1868 Proc. p. XXXVII.

Eucharissa Natalica Westwood, Thesaur. entom. Oxon. 1874 p. 144 n. 2.

speciosa Westw. — Afr.: Cap.

Eucharissa speciosa (Klug) Westwood, Trans. Entom. Soc. London 1868 Proc. p. XXXVII.

Eucharissa speciosa Westwood, Thesaur. entom. Oxon. 1874 p. 143 n. 1; T. 27 F. 3.

stigmatica Westw. — ♂ — Afr.: Caffraria.

Eucharissa stigmatica Westwood, Thesaur. entom. Oxon. 1874 p. 144 n. 3, ♂; T. 27 F. 2.

34. *Subfam. Eucharidinae*

Förster, Hymen. Stud. II. 1856 p. 19, 22 & 42.

DESTEFANIA

Dalla Torre, Wien. entom. Zeitg. 1897 p. 88.

De Stefani-Perez, nom. propr.

synon.: Sternodes Destefani, Natural. Sicil. X. 1891 p. 118 [non Fischer 1837].*στεφνον, pectus.**pusateri* (Dest.) D. T. — ♀ — Eur.: Sicilia.

Sternodes Pusateri Destefani, Natural. Sicil. X. 1891 p. 118, ♀.

Destefania Pusateri m.

EUCCHARIS

Latreille, Hist. nat. Crust. & Insect. III. 1802 p. 210 n. 346.

*εὐχαρῖς, gratus.**synon.*: Cynips Fabricius, Villers, Gmelin, Olivier.*adscendens* Fabr. — ♀ ♂ — Eur. fere tota.

Cynips adscendens Fabricius, Mant. Insect. I. 1787 p. 251 n. 1.

Cynips adscendens Villers, C. Linnæi Entom. III. 1789 p. 77 n. 26.

Cynips adscendens Gmelin, Linné: Syst. nat. Ed. 13^a I. 5. 1790 p. 2649 n. 20.

Cynips adscendens Olivier, Encycl. méthod. Insect. V. 1790 p. 780 n. 1.

- Cynips adscendens* Fabricius, Entom. system. II. 1793 p. 100 n. 1.
Eucharis adscendens Fabricius, Syst. Piez. 1804 p. 157 n. 1.
Eucharis adscendens Latreille, Hist. nat. Crust. & Insect. XIII. 1805 p. 210.
Eucharis adscendens Panzer, Faun. Insect. German. VIII. 1805 P. 88 T. 10.
Eucharis adscendens Panzer, Krit. Revis. II. 1806 p. 96.
Chalcis adscendens Jurine, Nouv. méth. class. Hymén. 1807 p. 315, ♀ ♂.
Eucharis adscendens Klug, Magaz. Ges. naturf. Fr. Berlin I. 1807 p. 76 n. 25.
Eucharis adscendens Latreille, Gen. Crust. & Insect. IV. 1809 p. 20.
Eucharis adscendens Spinola, Ann. mus. hist. nat. XVII. 1811 p. 149.
Eucharis adscendens Lamarck, Hist. nat. anim. s. vert. IV. 1817 p. 160 n. 1.
Eucharis adscendens Nees, Hymen. Ichneum. affini. Monogr. II. 1834 p. 266 n. 1.
Eucharis adscendens Lamarck, Hist. nat. anim. s. vert. Ed. 2^a IV. 1835 p. 370 n. 1.
Eucharis adscendens Blanchard, Hist. nat. Insect. III. 1840 p. 257 n. 1.
Eucharis adscendens Westwood, Introd. mod. Classif. Insect. II. 1840 Synops. p. 5.
Eucharis adscendens Stephens, Illustr. Brit. Entom. Suppl. 1846 p. 5; T. 44 F. 1.
Eucharis adscendens Taschenberg, Hymen. Deutschl. 1866 p. 111, ♀ ♂.
Eucharis adscendens Westwood, Thesaur. entom. Oxon. 1874 p. 151; T. 28 F. 12.
Eucharis adscendens W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. 1886 p. 11.
aeginctus Walk. — ♂ — As.: Philippines.
Eucharis Aeginctus Walker, List Hymen. Brit. Mus. Chalcid. I. 1846 p. 86, ♂.
convergens Walk. — ♂ — As.: Ceylon.
Eucharis convergens Walker, Ann. & Mag. Nat. Hist. (3) VI. 1860 p. 358, ♂.
delicatula Walk. — ♂ — Austr.
Eucharis delicatula Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 377, ♂.
democles Walk. — ♂ — Austr.: New South Wales.
Eucharis Democles Walker, Monogr. Chalcid. II. 1839 p. 15 n. 9, ♂.
dicerodera Spin. — ♂ — Am.: Brasilia.
Eucharis dicerodera Spinola, Mem. acad. sc. Torino (2) XIII. 1851 p. 43 n. 24, ♂.
eribotes Walk. — ♀ ♂ — Austr.: Tasmania, New South Wales.
Eucharis Eribotes Walker, Monogr. Chalcid. II. 1839 p. 14 n. 7, ♀ ♂.
fausta Walk. — ♂ — Austr.: Tasmania.
Eucharis fausta Walker, Monogr. Chalcid. II. 1839 p. 10 n. 2, ♂.
flabellata Fabr. — Am. mer.
Eucharis flabellata Fabricius, Syst. Piez. 1804 p. 158 n. 5.
gibbosa Prov. — ♀ — Am.: Canada.
Eucharis gibbosa Provancher, Natural. Canad. XII. 1881 p. 292, ♀.
Eucharis gibbosa Provancher, Faun. entom. Canada. Hymén. 1883 p. 571, ♀.
implexa Walk. — ♀ — Austr.: Tasmania.
Eucharis implexa Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 377, ♀.
kollari Först. — ♀ ♂ — Eur.: Austria inf.
Eucharis Kollari Förster, Verh. naturh. Ver. preuss. Rheinl. XVI. 1859 p. 91 n. 25, ♀.
larymna Walk. — ♀ ♂ — Austr.
Eucharis Larymna Walker, List Hymen. Brit. Mus. Chalcid. I. 1846 p. 86, ♀ ♂.
myrmiciae For.¹⁾ — ♀ ♂ — As.: India.
Eucharis (Chalcis) myrmiciae (Cameron) Forel, Ann. soc. entom. Belge XX. 1890 C.R. p. VIII.
Eucharis myrmiciae Cameron, Mem. & Proc. Manchester Soc. (4) IV. 1891 p. 187, ♀ ♂; T. 17.
piceicornis Walk. — ♂ — Austr.
Eucharis piceicornis Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 374, ♂.

¹⁾ *Hymen.*: *Myrmecia forficata* Fabr. (Cameron).

- punctata* Först. — ♀ — Eur. mer.
Eucharis punctata Förster, Verh. naturh. Ver. preuss. Rheinl. XVI. 1859 p. 89 n. 24, ♀.
pyttalus Walk. — Austr.
Eucharis Pyttalus Walker, List Hymen. Brit. Mus. Chalcid. I. 1846 p. 87.
rufiventris Walk. — ♀ — Austr.
Eucharis rufiventris Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 378, ♀.
smaragdina Walk. — ♂ — Austr.: Tasmania.
Eucharis smaragdina Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 376, ♂.
theocles Walk. — ♀ — Austr.: New South Wales.
Eucharis Theocles Walker, Monogr. Chalcid. II. 1839 p. 11 n. 4, ♀.
testacea Motsch. — As.: Ceylon.
Eucharis? testacea Motschulsky, Bull. soc. natural. Moscou XXXVI. P. 3. 1863 p. 44.
valgius Walk. — ♂ — Austr.: New South Wales.
Eucharis Valgius Walker, Monogr. Chalcid. II. 1839 p. 11 n. 3, ♂.
xeniades Walk. — ♂ — Austr.: New South Wales.
Eucharis Xeniaades Walker, Monogr. Chalcid. II. 1839 p. 15 n. 8, ♂.

ORASEMA

Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 105.

Etymol. obscura.

Synon.: *Eucharis* Fabricius, Walker.

Schizaspidia Westwood.

cameronii How. — ♀ — Am.: Grenada.

Orasema Cameroni Howard, Journ. Linn. Soc. London. Zool. XXVI. 1896 p. 133, ♀.

festiva (Fabr.) Kby. — Am. mer.

Eucharis festiva Fabricius, Syst. Piez. 1804 p. 157 n. 2.

Orasema festiva W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 29.

maculata (Westw.) Kby. — Am.: Brasilia.

Schizaspidia? maculata Westwood, Thesaur. entom. Oxon. 1874 p. 153; T. 28 F. 1.

Orasema maculata W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 29.

minuta Ashm. — ♂ — Am.: Florida.

Orasema minuta Ashmead, Entom. Amer. III. 1888 p. 188, ♂.

minutissima How. — ♀ ♂ — Am.: St. Vincent.

Orasema minutissima Howard, Journ. Linn. Soc. London. Zool. XXV. 1894 p. 84, ♀ ♂.

occidentalis Ashm. — ♀ — Am.: California.

Orasema occidentalis Ashmead, Proc. Entom. Soc. Washington II. 1893 p. 355 n. 3, ♀.

rapo (Walk.) Kby. — ♀ — Am.: Brasilia.

Eucharis Rapo Walker, Monogr. Chalcid. II. 1839 p. 66 n. 2, ♀.

Orasema Rapo W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 29.

sulthill How. — ♀ — Am.: Grenada.

Orasema Smithi Howard, Journ. Linn. Soc. London. Zool. XXVI. 1896 p. 134, ♀.

stramineipes Cam. — Am.: Panama. [F. 20 & T. 6 F. 18.]

Orasema stramineipes Cameron, Biol. Centr.-Amer. P. 31. 1881 Hymen. I. p. 105 n. 1, ♀; T. 5

Orasema flavipes Cameron, Biol. Centr.-Amer. P. 31. 1881 Hymen. I. p. 105 n. 1; T. 5 F. 20.

Orasema flavipes W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 29.

violacea Ashm. — ♂ — Am.: Florida.

Orasema violacea Ashmead, Entom. Amer. III. 1888 p. 187, ♂.

viridis Ashm. — ♀ — Am.: Mexico.

Orasema viridis Ashmead, Proc. Californ. Acad. Sc. V. 1895 p. 553 n. 33, ♀.

TRICORYNA

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 29 n. 3.

τρῆις, tres; *κορύνη*, clavus.

Synon.: *Eucharis* Walker.

jello (Walk.) Kby. — ♂ — Austr.: Tasmania.

Eucharis Jello Walker, Monogr. Chalcid. II. 1839 p. 12 n. 5, ♂.

Eucharis Jello Walker, Entomologist 1841; Tab. P Fig. 3.

Eucharis Jello Walker, Notes on Chalcid. P. 4. 1871 p. 66; Fig. 3.

Eucharis Jello Walker, Entomologist VI. 1872 p. 65; Fig.

Tricoryna Jello W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 29, ♂

METAGEA

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 30 n. 4.

Etymol. obscura.

Synon.: *Eucharis* Walker.

schwarzii Ashm. — ♀ ♂ — Am.: Columbia, Maryland.

Metagea Schwarzii Ashmead, Proc. Entom. Soc. Washington II. 1893 p. 356 n. 2, ♀ ♂

zalates (Walk.) Kby. — ♀ ♂ — Austr.: New South Wales.

Eucharis Zalates Walker, Monogr. Chalcid. II. 1839 p. 13 n. 6, ♀ ♂.

Eucharis Zalates Walker, Entomologist 1841; Tab. P Fig. 4.

Eucharis Zalates Westwood, Thesaur. entom. Oxon 1874 p. 151; T. 28 F. 15.

Metagea Zalates W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 30, ♀

PSILOGASTER

Blanchard, Hist. nat. Insect. III. 1840 p. 260.

ψῖλος, nudus; *γαστήρ*, venter.

cuprea Blanch. — Afr.: Aegyptus.

Psilogaster cupreus Blanchard, Hist. nat. Insect. III. 1840 p. 260.

Psilogaster cupreus W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 30

pallidipes Brullé — ♀ ♂ — Austr.: Tasmania.

[& 2 C

Psilogaster pallipes Brullé, Hist. nat. Insect. Hymén. IV. 1846 p. 574 n. 1, ♀ ♂; T. 39 F. 11

Psilogaster pallipes Desmarest, Chenu: Encycl. hist. nat. Anelles 1860 p. 161; Fig. 12

CHALCURA

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 30 n. 6.

χαλκός, aes; *οὐρά*, cauda.

Synon.: *Eucharis* Walker, Westwood.

americana How. — ♀ — Am.: St. Vincent.

Chalcura Americana Howard, Journ. Linn. Soc. London. Zool. XXV. 1894 p. 85, ♀

bedelii Cam.¹⁾ — ♀ ♂ — Afr.: Algeria.

Chalcura Bedeli Cameron, Mem. & Proc. Manchester Soc. (4) IV. 1891 p. 188, ♀ ♂; T. 1 F. 6 & 7

californica Ashm. — ♂ — Am.: California.

Chalcura Californica Ashmead, Proc. Entom. Soc. Washington II. 1892 p. 356 n. 2, ♂

¹⁾ *Hymen.*: *Myrmecocistus* viaticus (Cameron).

deprivata (Walk.) Kby. — ♀ ♂ — As.: Ceylon.

Eucharis deprivata Walker, Ann. & Mag. Nat. Hist. (3) VI. 1860 p. 359, ♂.

Eucharis deprivata Westwood, Thesaur. entom. Oxon. 1874 p. 155, ♀ ♂; T. 28 F. 6.

Chalcura deprivata W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 30.

RHIPIPALLUS

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31 n. 7.

ῥίπις, flabellum; πῖλλον, jacere.

Synon.: *Eucharis* Walker, Westwood.

Stilbula Walker.

cameronii Kby. — ♂ — Austr.? As.: Celebes? [n. 8, ♂; T. 1 F. 2.]

Rhipipallus Cameroni W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116: 1886 p. 37

turneri Kby. — Austr.

Rhipipallus Turneri W. F. Kirby, Ann. & Mag. Nat. Hist. (6) XIV. 1894 p. 47.

volusus (Walk.) Kby. — ♂ — Austr.

Eucharis Volusus Walker, Monogr. Chalcid. II. 1839 p. 9 n. 1, ♂.

Eucharis Volusus Walker, Entomologist 1841; Tab. P Fig. 1.

Stilbula Volusus Walker, Notes on Chalcid. P. 4. 1871 p. 65; Fig. 1.

Stilbula Volusus Walker, Entomologist VI. 1872 p. 65; Fig.

Eucharis Volusus Westwood, Thesaur. entom. Oxon. 1874 p. 151; T. 28 F. 14.

Rhipipallus Volusus W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31.

STILBULA

Spinola, Ann. mus. hist. nat. XVII. 1811 p. 150 n. 17.

Stilbum, nom. propr. Hymen.

Synon.: *Eucharis* Latreille, Nees, Blanchard.

Ichneumon Rossi.

cynipiformis (Rossi) Spin. — ♀ — Eur. mcr.

Ichneumon cynipiformis Rossi, Mant. Insect. 1792 p. 125 n. 275.

Eucharis cynipiformis Latreille, Gen. Crust. & Insect. IV. 1809 p. 21.

Stilbula cynipiformis Spinola, Ann. mus. hist. nat. XVII. 1811 p. 150.

Eucharis cynipiformis Nees, Hymen. Ichneum. affin. Monogr. II. 1834 p. 267 n. 2.

Eucharis cynipiformis Blanchard, Hist. nat. Insect. III. 1840 p. 257 n. 2.

Stilbula cynipiformis W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31.

grenadensis How. — ♀ — Am.: Grenada.

Stilbula Grenadensis Howard, Journ. Linn. Soc. London. Zool. XXVI. 1896 p. 133, ♀.

montana Ashm. — ♀ — Am.: Colorado.

Stilbula montanus Ashmead, Bull. Colorado Biol. Assoc. I. 1890 p. 24, ♀.

peduncularis Westw. — ♀ ♂ — Austr.

Stilbula peduncularis Westwood, Thesaur. entom. Oxon. 1874 p. 155, ♀ ♂; T. 28 F. 7.

SCHIZASPIDIA

Westwood, Proc. Zool. Soc. London III. 1835 p. 69.

σχίζω, findo; ἄσπις, scutum.

caffa Westw. — Afr.: Caffraria.

Schizaspidia Caffra Westwood, Thesaur. entom. Oxon. 1874 p. 152 n. 4; T. 28 F. 12.

cyanca Walk. — ♂ — As.: Amboina.

Schizaspidia cyanca Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 386, ♂.

furcifera Westw. — As.: Bengal.

Schizaspidia furcifera Westwood, Proc. Zool. Soc. London III. 1835 p. 69.

Schizaspidia furcifera Westwood, Thesaur. entom. Oxon. 1874 p. 151 n. 1. ♀?; T. 28 F. 1

Schizaspidia furcifera W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31.

murrayi Kby. — ♂ — Austr.: Tongatabu.

Schizaspidia Murrayi W. F. Kirby, Ann. & Mag. Nat. Hist. (5) XIII. 1884 p. 403 n. 2. ♂.

Schizaspidia Murrayi W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 37. ♂.

pretendens Walk. — ♂ — Am.: Brasilia. [T. 1 F. 1]

Schizaspidia pretendens Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 385. ♂.

rudis Westw. — Austr.

Schizaspidia rudis Westwood, Thesaur. entom. Oxon. 1874 p. 152 n. 2; T. 28 F. 5.

TETRAMELIA

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31 n. 10.

τέτρα, quattuor; *μελία*, fraxinus.

Synon.: *Schizaspidia* Walker.

meridionalis Kby. — ♀ — Am.: Brasilia.

Tetramelia meridionalis W. F. Kirby, Ann. & Mag. Nat. Hist. (6) IV. 1889 p. 144. ♀.

plagiata (Walk.) Kby. — ♂ — Am.: Brasilia.

Schizaspidia plagiata Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 385. ♂.

Schizaspidia plagiata Westwood, Thesaur. entom. Oxon. 1874 p. 152 n. 3; T. 28 F. 11

Tetramelia plagiata W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31.

LOPHYROCERA

Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 103.

Lophyrus, nom. propr. Hymen.; *κέρας*, cornu.

apicalis Ashm. — ♀ — Am.: California.

Lophyrocera apicalis Ashmead, Proc. Entom. Soc. Washington II. 1893 p. 357 n. 3. ♀.

floridana Ashm. — ♂ — Am.: Fla.

Lophyrocera floridana Ashmead, Entom. Amer. III. 1888 p. 187. ♂.

niromaculata Cam. — ♂ — Am.: Arizona, Nicaragua. [T. 5 F. 19]

Lophyrocera niromaculata Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 104 n. 2. ♂.

Lophyrocera niromaculata W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886

stramineipes Cam. — ♀ — Am.: Panama. [p. 32 a. 1]

Lophyrocera stramineipes Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 103 n. 1. ♀.

[T. 5 F. 19]

Lophyrocera stramineipes W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886

[p. 32 a. 1]

KAPALA

Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 102.

Etymol. obscura.

Synon.: *Chirocera* Desmarest.

Chirocerus Brullé.

Eucharis Fabricius, Latreille, Lamarck, Walker.

Schizaspidia Walker.

Thoracantha Westwood, Walker.

furcata (Fabr.) Cam. — ♀ ♂ — Am.: Brasilia, Grenada; Afr.: Sierra Leone.

Eucharis furcata Fabricius, Syst. Piez. 1804 p. 158 n. 4. ♀.

Eucharis flabellatus Fabricius, Syst. Piez. 1804 p. 158 n. 5. ♂.

- Eucharis furcata* Latreille, Gen. Crust. & Insect. III. 1807 p. 21.
Eucharis furcata Lamarek, Hist. nat. anim. s. vert. IV. 1817 p. 160 n. 2.
Eucharis furcata Lamarek, Hist. nat. anim. s. vert. Ed. 2^a IV. 1835 p. 370 n. 2.
Eucharis furcata Walker, Monogr. Chalcid. II. 1839 p. 65 n. 1, ♀.
Thoracantha furcata Walker, Entomologist 1841; Tab. P Fig. 2.
Chirocerus furcatus Brullé, Hist. nat. Insect. Hymén. IV. 1846 p. 571; T. 38 F. 5.
Chirocerus furcatus Lucas, Sagra: Hist. fis. etc. Cuba VII. 1856 p. 762.
Chirocera furcata Desmarest, Chenu: Encycl. hist. nat. Annelles 1860 p. 161; Fig. 141.
Schizaspidia furcata Walker, Notes on Chalcid. P. 4. 1871 p. 66; Fig. 2.
Schizaspidia furcata Walker, Entomologist. VI. 1872 p. 85; Fig.
Kapala furcata Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 103 n. 1, ♀; T. 5 F. 17.
Kapala furcata W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 32.
terminalis Ashm. — ♂ — Am.: Cuba.
Kapala terminalis Ashmead, Proc. Entom. Soc. Washington II. 1892 p. 358, ♂.

THORACANTHA

Latreille, Fam. nat. règn. anim. 1825 p. ?.

θώραξ, thorax; ἄκανθα, aculeus.

Synon.: *Galecaria* Brullé, Hist. nat. Insect. Hymén. IV. 1846 p. 527.
 Etymol. obscura.

Suby.: *Scutellista* Motschulsky, Étud. entomol. VIII. 1859 p. 172.
 scutellum.

alta Walk. — ♀ — Am.

Thoracantha alta Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 383, ♀.

anchurus Walk. — Am.: Brasilia.

Thoracantha Anchurus Walker, List Hymen. Brit. Mus. Chalcid. I. 1846 p. 88.

atrata Walk. — ♂ — Patria?

Thoracantha atrata Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 383, ♂.

batesella Westw. — Am.: Brasilia.

Thoracantha Batesella Westwood, Thesaur. entom. Oxon. 1874 p. 154 n. 5; T. 28 F. 8.

bella Först. — ♀ — Eur. mer.

Thoracantha bella Förster, Verh. naturh. Ver. preuss. Rheinl. XVI. 1859 p. 92 n. 26, ♀.

cyanea (Motsch.) D. T.¹⁾ — ♀ ♂ — As.: Ceylon.

Scutellista cyanea Motschulsky, Étud. entom. VIII. 1859 p. 172, ♀ ♂; T. 1 F. 17.

Thoracantha cyanea m.

cynipsea Walk. — ♀ ♂ — Am.: Brasilia.

Thoracantha cynipsea Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 379, ♀ ♂.

floridana Ashm. — ♀ ♂ — Am.: Florida.

Thoracantha Floridana Ashmead, Entom. Amer. I. 1885 p. 96, ♂.

Thoracantha Floridana Ashmead, Trans. Amer. Entom. Soc. XII. 1885 Proc. p. XI n. 5, ♀ ♂.

luxagens Walk. — ♀ — Am.: Brasilia.

Thoracantha inexagens Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 381, ♀.

latreillei Guér. — ♀ — Am.: Brasilia.

Thoracantha Latreillei Guérin, Iconogr. règn. anim. VII. Insect. 1829-45 p. 415 n. 1; T. 67 F. 8.

Thoracantha Latreillei vel coleopteroides (Waterhouse: Westwood, Trans. Entom. Soc.

[London II. P. 3. 1839 p. 196; T. 17 F. 3.

Thoracantha Latreillei Blanchard, Hist. nat. Insect. III. 1840 p. 257 n. 1.

¹⁾ *Rhynch.*: *Lecanium coffeae* (Motschulsky).

- Thoracantha Latreillei Walker, Ann. & Mag. Nat. Hist. XII. 1843 p. 45, ♀.
 Galearia violacea Brullé, Hist. nat. Insect. Hymén. IV. 1846 p. 572 n. 1; T. 38 F. 6.
 Thoracantha Latreillei W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 31
 reflexa Walk. — ♀ — Am.: Brasilia.
 Thoracantha reflexa Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 382, ♀.
 romandii Guér. — Am.: Columbia.
 Thoracantha Romandii Guérin, Iconogr. règn. anim. VII. Insect. 1845 p. 415 n. 2.
 striatissima Walk. — ♀ ♂ — Am.: Brasilia.
 Thoracantha striatissima Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 380, ♂♂.

ACROSTELA

Shipp, Entomologist XXVII. 1894 p. 188.

ἄκροσ, summus; στῆλη, columna.

Synon.: Thoracantha Latreille.

apta (Walk.) Shipp — ♂ — Am.: Brasilia.

Thoracantha apta Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 384, ♂.

Thoracantha apta Westwood, Thesaur. entom. Oxon. 1874 p. 153 n. 1; T. 28 F. 3.

Acrostela apta Shipp, Entomologist XXVII. 1894 p. 188.

DILOCANTHA

Shipp, Entomologist XXVII. 1894 p. 188.

Etymol. obscura.

Synon.: Thoracantha Walker, Westwood.

f/avicornis (Walk.) Shipp — ♀ — Am.: Brasilia.

Thoracantha flavicornis Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 382, ♀.

Thoracantha flavicornis Westwood, Thesaur. entom. Oxon. 1874 p. 153 n. 2; T. 28 F. 4.

Dilocantha flavicornis Shipp, Entomologist XXVII. 1894 p. 188.

ISOMERALA

Shipp, Entomologist XXVII. 1894 p. 188.

ἴσος, aequus; μηριαῖος, femoralis.

Synon.: Thoracantha Westwood.

coronata (Westw.) Shipp — Am.: Brasilia.

Thoracantha coronata Westwood, Thesaur. entom. Oxon. 1874 p. 154 n. 4; T. 28 F. 10.

Isomerala coronata Shipp, Entomologist XXVII. 1894 p. 188.

LAETOCANTHA

Shipp, Entomologist XXVII. 1894 p. 188.

λαῖτος, publicus; ἄκανθα, aculeus.

Synon.: Thoracantha Walker.

nasua (Walk.) Shipp — ♂ — As.: Philippines Ins.

Thoracantha Nasua Walker, List Hymen. Brit. Mus. Chalcid. I. 1846 p. 88, ♂.

Laetocantha Nasua Shipp, Entomologist XXVII. 1894 p. 188.

LASIONYCHUS

Shipp, Entomologist XXVII. 1894 p. 188.

λάσιος, pubescens; ὄνυξ, unguis.

Synon.: Chalcis Blanchard.

Thoracantha Westwood, Blanchard.

flabellatus (Westw.) Shipp — ♀ ♂ — Am.: Brasilia.

Thoracantha flabellata Westwood, Proc. Zool. Soc. London III. 1835 p. 52, ♂ [non Fabricius].

Thoracantha aculeata Westwood, Proc. Zool. Soc. London III. 1835 p. ? ♀.

Chalcis (Thoracantha) aculeata Blanchard, Cuvier: Règne anim. Ed. 3^e Insect. II. 1849; T. 113

Thoracantha aculeata Westwood, Thesaur. entom. Oxon. 1874 p. 154 n. 3; T. 28 F. 9. [F. 8.

Lasionychus flabellatus Shipp, Entomologist XXVII. 1894 p. 188.

LIRATA

Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 102.

Etymol. obscura.

Synon.: Schizaspidia Cameron.

Thoracantha Walker.

luteogaster Cam. — Am.: Mexico.

Lirata luteogaster Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 102 n. 1; T. 5 F. 16.

Schizaspidia luteogaster Cameron, Biol. Centr.-Amer. P. 31. 1884 Hymen. I. p. 102; T. 5 F. 16.

Lirata luteogaster W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 33.

pallescens (Walk.) Shipp — ♂ — Am.: Brasilia.

Thoracantha pallescens Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 380. ♂.

Lirata pallescens Shipp, Entomologist XXVII. 1894 p. 188.

surgens (Walk.) Shipp — ♂ — Am.: Brasilia.

Thoracantha surgens Walker, Trans. Entom. Soc. London (3) I. P. 4. 1862 p. 384. ♂.

Lirata surgens Shipp, Entomologist 1894 p. 188.

UROMELIA

W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 33 n. 15.

ὄρα, cauda; μελία, terra melica.

Synon.: Thoracantha Perty, Blanchard.

striata (Perty) Kby. — Am.: Brasilia.

Thoracantha striata Perty, Delect. anim. artic. Brasil. 1833 p. 134; T. 28 F. 15 & 16.

Thoracantha striata Blanchard, Hist. nat. Insect. III. 1840 p. 257 n. 2.

Uromelia striata W. F. Kirby, Journ. Linn. Soc. London. Zool. XX. P. 116. 1886 p. 33.

35. Subfam. Chalcidinae

Förster, Hymen. Stud. II. 1856 p. 18, 21 & 29.

DIRRHINUS

Dalman, Svensk. Vet.-Akad. Handl. XXXIX. 1818 p. 75 n. 3.

δῆς, bis; ῥῖν, nasus.

anthracia Walk. — As.: Philippines.

Dirrhinus Anthracia Walker, List Hymen. Brit. Mus. Chalcid. I. 1846 p. 85.