

Micromelius Sutanus, Rem. *Aneno-niger, cyaneo-varius, antennae fulvo-*
pedes fulvi, ales fuscæ.

Corpus xeno-nigrum, su-

卷之三

Genus *LILARS*, *Holiday MSS.*

Corpus aeneo-nigrum, subbus cyaneo-nigrum, breve, latum, convexum, parvum nitens, scutissime squamatum, parce hirtum: caput transversum, breve, thorace tatus; vertex tatus; fons impressus, abrupte declivis: oculi picte, mediocres, non extantes: ocelli vertice triangulum flingentes; medius perparvum antepositus: antennae fulvae, ad os inserentes, basi graciles lineares: os picuum: thorax ovatus: prothorax brevissimum: mesothoracis scutum longitudine latius, parapsides scuto fere in unum confusae: parapetra et epimera magna; scutellum subtriangulariforme, postice latius: metathorax brevis, declivis, postice angustius: petiolis brevissimum: abdomen subrotundum, tenuum, nitens, laeve, supra planum, subbus carinatum, thorace paulo latius et multo brevius: pedes fulvi, simplices, subrquales: ate fusce, mediocres; squamulae picore; nervi fuscii; nervus humeralis ulnari duplo longior, rami diuersi ulnari brevior cubitali vix longior, cubitalis sat longus; stigma medio-dioce. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{4}$.)

VIII.—*Descriptions of Chaileidites discovered in St. Vincent's Isle by the Rev. Lansdown Guilding.* By FRANCIS WALKER, Esq.,

Decatoma Oretilia, Mas et Fem. *Fulvo-flava ferrugineo-varia*, antennæ fulvae, pedes flavi, alici impilie.

Corpus convexum, sublineare, fulvo-flavum, nitens, subtilissime punctatum, ante hirtum : caput transversum, breve, thorace paullo latius ; vertex latus ; tons impressa, abrupte declivis : oculi ruti, medocres, non extantes : annulus fulvus, elongatus, thoraco non transverso, rotundatus. Tarsus longus, levigatus.

lutescentes; clava fusiformis, acuminate, articulo 9° multo laetior et plus duplo longior; thorax ovatus: prothorax maximus, transversus, subquadatus, postice ferrugineo-maculatus: mesothoracis scutum brevissimum: parapsidum sutura vix conspicue: scutellum subrotundum: discus ferrugineus: metathorax transversus, brevis: petiolus brevis: abdomen ovatum, tave: *mari*: thoracs longitudine, *fem.* thorace paulo longius subtus carinatum apice acuminatum fuscum: vitta per dorsum interrupta fuscata. Pedes pallide flavi, simplices, subaequales: alae lumbipide: squamulae fulvae: nervi fulvi: nervi humerulari ulnari fore duplo longior: radialis ulnari non brevir capitulo plus duplo longior; stigma minutum. (Corps long. lin. 1; alar. lin. 1½.)

Pteronotus Helice, Tem. *Aeneo-viridis*, antennæ fulvæ, pedes flavi, alæ
impide.

Corpus convexum, aeneo-virende, nitens, scissime squameum, parce hirtum: caput transversum, breve, thorace latius; vertex latus; frontis impressa, abrupte declivis: oculi rufi, mediocre non extantes: antennae fulvæ, extorris, crassiores, thorace vix longiores; articulus 1^{us} longus, gracilis; 2^{us} cylindricus; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, usque ad 10 mm latentes; clava longicirca, acuminata, articulo 10° duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudo latius; parastidium suture vix conspicue: scutellum subconicum, angustius: metathorax obconicus, nictocoris, declivis: petiolis brevissimis: abdomen fusiforme, levè supra depresso, subito curvatum, apice acuminatum, thorace longius et angustius: pedes tate flavi, simplices, subæquales: alæ limpidae; squamulae fulvæ; nervi flavi; nervus humeralis

Lasioglossum (Halictus) MSS., Fem. *Cyprea, antennae fuscæ flavo-piceæ, pedes fuscæ, proboscis fuscæ-nigrae.*

C. c. capreum; viridi-varium: oculi et ocelli rufi; antennae fuscae, flavo-
-luteae, apice flavae; articulus 1^{us} flavus: ovipunctus vagine fulvæ, apice
-luteo-fuscæ; pretex. flavi; ungues et pulvilli fusi: aleæ limpidae; proala fusco-
-lutea, ang. superiore fulvae; nervi fulvi. (Corp. long. lin. 1½; alar. lin. 2¾.)

卷之三

Thysan angustus; transversus: mesothoracis scutum impessum, longitudo scutellae; parastigma sutura remota, bene determinata, postice approximata; scutulum magnum, transversum, bisulcatum; metathorax magnum, tricuspidatum; petiolus vix ullus; abdomen longi-ovatum, thorace paullo rotundatum; segmentum 1^{um} magnum, 2^{um} et sequentia brevia: 3^{um} et 4^{um} longissimum; vaginæ compone triplo longiores; propedum coxa longior; femora crassa; tibiae late: mesopodes graciles; coxa parvæ: scutellum exæ sat magne; femora crassa; tibiae subarcuatae: tarsorum tenuia; latæ: cunctæ, 5^{um} et 6^{um} paullo longior; ungues et pulvilli minuti; aenescens; nervus humeralis longus, ulnaris brevissimus, radialis longus.

Theridion P. in. *Virilius-nodus*, abdomen cupreum, antennæ nigrae, pedes albi tenuide.

Coccinellidium: oculi et ocelli rufi: antennae nigre: abdomen curvatum: vittulae fuscis: vagina picea: pedes fulvi: immixta et rotundata.

Pterostichus Garganus, Mas. *Furcatus, abdomen cupreum, antennae fulve,*
petiolaris, ala albo-limpida.

THE ANNALS

AND

MAGAZINE OF NATURAL HISTORY,

INCLUDING

ZOOLOGY, BOTANY, AND GEOLOGY.

(BEING A CONTINUATION OF THE 'MAGAZINE OF BOTANY AND ZOOLOGY,' AND OF
LOUDON AND CHARLESWORTH'S 'MAGAZINE OF NATURAL HISTORY.')

CONDUCTED BY

SIR W. JARDINE, BART., F.L.S.—P. J. SELBY, Esq., F.L.S.,

GEORGE JOHNSTON, M.D.,

CHARLES C. BABINGTON, Esq., M.A., F.L.S., F.G.S.,

J. H. BALFOUR, M.D., Reg. Prof. Bot. Glasg.,

AND

RICHARD TAYLOR, F.L.S., F.G.S.

VOL. XII.

LONDON:

PRINTED AND PUBLISHED BY R. AND J. E. TAYLOR.

SOLD BY S. HIGLEY; SIMPKIN AND MARSHALL; SHERWOOD AND CO.; W. WOOD,
TAVISTOCK STREET; BAUILLIÈRE, REGENT STREET, AND PARIS;
LIZARS, AND MACLACHLAN AND STEWART, EDINBURGH;
CURRY, DUBLIN: AND ASHER, BERLIN.

1843.